

#10

Paper March 24 1825

W. S. H.

Inaugural Essay  
on  
Rheumatismus acutus

by  
John R. Wise of V.  
March 10<sup>th</sup> 1825.

Copy of the

1831

Shardour & Co  
of the same name

of the same name

of the same name

Rhe  
order  
to  
and  
by  
Hollow  
lains  
to the  
more  
It rec  
bring  
Two  
season  
time  
lost  
of the  
appear  
does not  
after

Rheumatismus Acutus. Class Pyrexia  
order Phlegmasia of Cullen, who defines  
it to be a disease from an internal  
and often evident cause, attended  
by Pyrexia, pains about the joints  
following the course of the muscles,  
fixing on the larger joints in preference  
to those of the feet and hands.  
increased by external heat.

It occurs most frequently in autumn and  
spring, when there are sudden vicissitudes  
of weather; but may happen at any  
season if those changes are for the  
time present.

Sometimes the pains take the precedence  
of the fever; but in other cases the fever  
appears first, and the local affection  
does not discover itself till a few days  
afterwards.

*[Faint, mostly illegible handwriting on the left page, possibly bleed-through from the reverse side.]*

The 1st  
from a  
shell -  
and ho  
and pa  
usual  
spirit,  
acting  
the ton  
of a br  
travel  
high -  
the co  
sedom  
pains a  
com more  
and P  
and co  
we sho



The symptoms are the same as of fever from any other cause. Such as cold chills. Succeeded by a full, frequent and hard pulse. flushings of the face and partial heats; aversion to food, general lassitude, and depression of spirits, more or less of soreness and aching over different parts of the body. The tongue becomes coated with fur of a brownish hue, the thirst insupportable bowels costive, urine scanty and high-coloured, depositing during the course of the disease a turbid sediment. In the course of a short time pains are felt in different parts: most commonly the larger joints - the Hips and Knees of the lower, and shoulders and elbows of the upper extremities. we should recollect, however, that

the same  
from  
the an  
The P  
very co  
during  
the Pa  
and we  
one join  
The lin  
seat of  
become  
smaller  
which  
and ab  
does not  
We are  
in the c  
time

the smaller joints are not entirely  
free from its attacks; more especially  
the ancles and wrists.

The Gout suffers an exacerbation  
very evening, which is most considerable  
during the night, at which time  
the Pains also are much aggravated,  
and exceedingly prone to shift from  
one joint to another.

The limb which is the immediate  
seat of the disease, sooner or later  
becomes affected with redness and  
swelling at the coming on of  
which symptoms, there is most generally  
an abatement of the pain: but this  
does not invariably happen.

We are told by Cullen, that early  
in the course of this disease there is  
some sweating, but that it is seldom

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

few or  
the bac  
Blood  
an inf  
Rheum a  
in this  
is sub  
Fishes  
that it  
the m  
by the  
says, tha  
himself  
catus i  
a case  
recurre  
of Mary  
after t  
to h  
0

few or copious, and seldom either relieves  
the pain or proves critical.

Blood drawn in this diseased state is  
an inflammatory or sily appearance.  
Rheumatism differs from common inflammation  
in this (viz) that it very rarely terminates  
in suppuration, or gangrene; some  
Authors have gone so far as to affirm  
that it never does terminate in  
this manner; but this is contradicted  
by the observation of Dr Good, who  
says, that, in one or two instances he  
himself had been witness to an  
extensive abscess. I have also seen  
a case of the same nature, which  
recurred to my preceptor Dr John Purnell  
of Maryland. Here, it became necessary  
after trying by every possible means  
to bring absorption of the pus  
about.

to man  
it might  
difficult  
the spe  
the wo  
and the  
great a  
relation  
best of  
it to m  
whilest  
in the  
tendency  
will to  
organs  
flowers  
which  
to feel  
from the

to make an opening through which  
it might be discharged. Considered  
difficulty was experienced in causing  
the opening thus made to cicatrize;  
this was however ultimately accomplished  
and the patient perfectly recovered.  
Great diversity of opinion exists  
relative to the most common  
seat of this disease; some supposing  
it to be seated in the blood vessels,  
whilst others hold it to be seated  
in the loose mucous and contiguous  
tendinous membranes. Others again  
will tell us, that the muscular  
organs are alone affected. Richat  
observes that "it is yet to be ascertained  
which is attacked the muscular  
or fibrous texture." we can account  
for this difference of opinion in

is to  
 for a  
 that  
 A firm  
 by Sec  
 causes  
 it is o  
 be dr  
 read o  
 The fo  
 more  
 when  
 causes  
 such  
 might  
 belong  
 of wh  
 the p



no other way, than by attributing  
it to the want of opportunities  
for ascertaining the morbid  
Anatomy of the disease.

A Hereditary structure is assumed  
by Scudamoro as one of the remote  
causes of this disease; but says "it  
is obvious that the inference can  
be drawn only from general  
reasonings and not from demonstrations."

The fact, that, some persons are  
more liable than others to an attack,  
when equally exposed to its exciting  
causes, would seem to favour  
such an opinion; though, this  
might be owing to some peculiar  
idiosyncrasy of Constitution independently  
of any hereditary predisposition.  
The most common remote cause,



is the application of cold to the whole, or part of the body, when unduly heated by exercise or other wise, wearing wet or moist clothes, or sleeping in damp sheets; strains, and spasms - indeed, tetanoid injuries of any kind may be enumerated as sometimes laying the foundation of this disease.

We now come to the consideration of the proximate cause, and here again we shall find great difference of opinion amongst writers. Sydenham supposes the proximate cause to consist in a deprivation of morbid matter to the limb. This hypothesis I believe has been long abandoned

by the  
as far  
notion  
bullet  
is the  
prode  
totema  
and a  
covered  
an in  
and s  
of it.  
thorn  
which  
proam  
I conce  
the in  
blood  
the a

by Physicians of the present day  
as partaking too much of the  
notions of the humoral Pathologists.  
Cullen believes, that, cold which  
is the most common remote cause  
produces constriction of the  
arterial vessels on the surface,  
and a phlogistic diathesis in the  
course of blood, which causes  
an increased impetus of blood  
and resistance to the free passage  
of it. Neither of the above mentioned  
theories comport with the views,  
which I entertain relative to the  
proximate cause of inflammation.  
I concur with Dr. Cullen in attributing  
the increased determinations of  
blood, which always happens, to  
the action of cold, which is the



most frequent remote cause, or the capillaries, by which they are debilitated or paralyzed in such a manner as to render them unable to propel their contents; a stagnation of blood takes place, because there is no power to keep it in motion. The vis a tergo, or force from behind continuing the same, produces an accumulation or undue determination of blood to the part thus debilitated.

The diseases with which Rheumatism is most liable to be confounded are Gout, Scurvy & Syphilis, and great difficulty is sometimes experienced in distinguishing between those affections. The following circumstances will generally serve

to the  
former  
more g  
rough  
and ly  
much  
of deske  
of the  
be a d  
which  
two a  
know  
of attac  
of rel  
with  
to be  
arrive  
of accu  
into t



to distinguish it from Gout, the former (i.e.) Rheumatism comes on more gradually, and has more regularly marked exacerbations, and less clear remissions. it is also much less connected with symptoms of dyspepsia, or disordered condition of the stomach; to which may be added the seldomness with which it attacks the joints of the toes and fingers, which we all know is the most common seat of attack of the Gout.

As relate to the other diseases with which Rheumatism is apt to be confounded, we may generally arrive at a tolerably just degree of accuracy, by a minute examination into the previous histories and habits

The first of these is the  
fact that the number of  
act of the  
organ-  
isms  
may be  
of the  
The  
Rheum  
the as  
that  
if prop  
those  
called  
been  
be a  
the as

of the  
has  
valis  
of act  
organ-  
isms  
number  
may be  
of the  
The  
Rheum  
the as  
that  
if prop  
those  
called  
been  
be a  
the as

of the patient.

This disease seldom terminates fatally unless by metastasis, or translocation of action to some vital part or organ. Such cases are exceedingly rare in comparison with the great number affected, and hence we may generally prognosticate favourably of the disease.

The duration of an attack of Rheumatism will depend in a great measure on the mode of practice that is pursued; seldom continuing (if properly managed) longer than three or four weeks.

Called to a case such as I have been treating of; we can scarcely be at a loss, as to the proper measures to be adopted. The prompt

*[Faint, mostly illegible handwriting in cursive script, likely bleed-through from the reverse side of the page.]*

extra at  
and to  
will be  
by the  
haas  
urgency  
Sydney  
Blum  
of blue  
who is  
the cu  
to the  
1875 to  
the pra  
is free  
the ha  
suborg  
appears  
blue d

detracting of blood is indispensable;  
and the propriety of its repetition  
will be clearly indicated, both  
by the good effects which it may  
have produced, and by the  
urgency of the subsequent symptoms.

Sydenham in his first essay on  
Rheumatism, advocates the propriety  
of bleeding as the chief remedy  
which should be employed in  
the cure. In a subsequent treatise  
on the Epidemic disease from the year  
1675 to 1680 he appears to regret  
the practice of taking away blood  
so freely, as he had before been in  
the habit of recommending. On a  
subsequent occasion however he still  
appears partial to the practice of  
bleeding.

Ringler  
 of the  
 success  
 by rep  
 by be  
 the ch  
 "The  
 in  
 he did  
 propo  
 and  
 to the  
 His f  
 being a  
 such to  
 found  
 the a  
 Rheum  
 deficien

Kingle, in his account of the disease  
of the army, also informs us of the  
successful treatment of acute Rheumatism  
by repeated bleedings.

By Cullen blood letting is considered  
the chief remedy of Acute Rheumatism  
"The blood ought to be drawn (says  
he, in large quantity, and the  
bleeding is to be repeated, in  
proportion to the frequency, fulness,  
and hardness of the pulse, and  
to the violence of the pain."

His fear, however of profuse bleedings  
being apt to produce chronic Rheumatism  
seems to me to be entirely without  
foundation; for I believe, that when  
the acute is changed into chronic  
Rheumatism, it is caused rather by  
deficiency, than excess of blood letting.

through  
is of the  
comedy  
of the  
bath  
in the  
render  
becomes  
diathese  
has been  
on the  
much  
the com  
as they  
vacant  
fact  
from m  
we can  
writer



enough has now been said to convince  
us of the vast importance of the  
remedy in the management  
of this disease.

Cathartics are also very serviceable  
in the early treatment of this disease.  
Under their operations, the circulation  
becomes moderated, and the phlogistic  
diathesis subdued. This practice  
has been objected to by Authors  
on the ground, that, they do as  
much harm by the motion which  
they compel the patient to make,  
as they do good by their power of  
evacuation. Whether this be the  
fact or not, I am unable to say  
from my own observations. but if  
we can credit the accounts of a  
writer of no ordinary character,

show  
as we  
a con  
to day  
pects  
can not  
is not  
for the  
slow or  
of Low  
What  
Practice  
It was  
Raygar  
the stor  
cleanse  
all a  
for let  
the e

when he says, that "in proportion  
as we pursue this practice upon  
a continued principle from day  
to day, do we obtain its good  
effects in acute Rheumatism" we  
cannot but conclude that this  
is not the case. The best Cathartics  
for the purpose are the Saline either  
alone or in union with an infusion  
of Senna and Manna.

What shall we say of the use of  
Emetics in the cure of this disease?  
It was the favourite practice of  
Haggarth to administer Emetics, until  
the Stomach, and bowels were completely  
cleansed, at the very onset of an  
attack, and we are told by  
Professor Chapman, that in that  
species of Rheumatism originating

the mica  
the att  
the m  
accu  
sur  
after  
reduc  
some  
recom  
on the  
natur  
surfac  
the m  
with  
an tim  
the hoo  
grain  
or two

in Miasmatic districts, when  
the attack is blundered with  
intermittent fever, and great  
accumulations of bile. He has  
seen it do good in <sup>one</sup> case.

After arterial action and general  
excitement, have been sufficiently  
reduced by the above mentioned  
remedies, we may then have  
recourse to diaphoretics; these  
are medicines which promote the  
natural discharge from the  
surface of the body - for this purpose  
the nitrous powder in combination  
with Calomel and tartarized  
antimony is an excellent formula.  
The hourly exhibition of one or two  
grains of Speacenanha, and ten  
or twelve of Nitre constitute also

an account  
must be  
given to the  
disease,  
has been  
me as well  
it is re-  
should  
cases, for  
than to  
diaphor-  
but as  
important  
our lin-  
the little  
as to the  
Park so  
one ser-

an excellent sudorific prescrip. we must not pass by unnoticed the Dover powder: this is well adapted to the secondary stages of this disease, when fibrile excitement has been completely subdued by measures formerly pointed out: it is recommended that the sweating should be kept up in obstinate cases, for not less on an average than twenty four hours. Many other diaphoretics might be mentioned: but as most of them are of minor importance, we shall not occupy our time in enumerating them. No little difference of opinion prevails as to the propriety of using the Peruvian Bark in acute Rheumatism, whilst one set of practitioners, most highly

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

...the ...  
...by a ...  
...ly to ...  
...ing the ...  
...ing M ...  
...ous of ...  
...rivers ...  
...Pappa ...  
...that ...  
...only w ...  
...philis ...  
...medicine ...  
...up can ...  
...hild ...  
...and with ...  
...ould ...  
...healing ...  
...to the Pa ...  
...through



extol it, there are others, who  
utterly condemn its use in the  
early or inflammatory stage.  
Among those who speak favourably  
of it, may be mentioned the  
names of Farrar, Fothergill and  
Saunders. The former of whom  
(i.e.) Farrar, goes so far as to  
say that "Bark in Rheumatism  
is only inferior to Mercury in  
syphilis." I have never seen the  
Medicine tried in the early  
or inflammatory stage, but  
should suppose it injurious,  
and with Professor Chapon and  
should say "that generally  
speaking it is best suited  
to the Convalescence, to recruit  
strength and confirm recovery."

*[The page contains faint, illegible handwriting, likely bleed-through from the reverse side.]*

typical  
on of good  
the disc  
when gen  
caused is  
usually  
be negle  
inflam  
after the  
advised  
points o  
the app  
the pa  
as a lo  
th am  
the lea  
and ap  
but M  
said s

Topical evacuations by cups and leeches  
are of great service in the cure of  
this disease, and may be employed  
when general bleeding from any  
cause is inadmissible, or as an  
auxiliary to it. They should never  
be neglected, if the pain, and  
inflammation are severe.

After the inflammation has been  
reduced by the remedies formerly  
pointed out, we shall find  
the application of a blister to  
the pained part of much benefit.  
As a local remedy the Satura-  
tramonium has been recommended.  
the leaves steeped in brandy,  
and applied to the limb, is the  
best mode of application, it is  
said that it sometimes affords

not refer  
him all  
over for  
commen  
mercury  
a down  
this m  
black  
long to  
mercury  
certa  
and pro  
touch  
but dis  
strange  
to buy  
of the P  
very other  
we are

Much relief.

When all the remedies above enumerated have failed to give relief, it is recommended to resort to the use of Mercury with a view to a salivation as a dernier alternative. The propriety of this measure has been disputed by Clarke, a writer on the diseases of long voyages. He says that "though Mercury is an other wise of the greatest importance, it never failed to aggravate and protract the complaint when it touched the mouth." In this I must disagree with him, having witnessed its beneficial effects so very frequently in the practice of the Pennsylvania Hospital, where every other remedy had failed. We are told by professor Chapman

the disease  
the mouth  
accomplish  
recurrence  
for a course  
than the  
nothing  
the various  
such as  
partial  
stiffness of  
allows  
series  
the prop  
consist-  
measures  
to streng  
a agric

That cases do sometimes occur, when the disease will not yield until the mouth is touched; and to accomplish a radical cure the mercurial impression must be sustained for a considerable period.

When the disease is removed and nothing, but its effects remain the various Tonics become proper: such as Bark, Mineral acids, Martial preparations &c. &c. The stiffness of the limb, which sometimes follows, is to be overcome by exercise, and friction.

The prophylactic management consist in the employment of those measures, which are calculated to strengthen the body, and fortify it against the impressions of those

the p  
d the  
Hannul  
during  
great in  
the diet  
simple, a  
as early  
fully, tips  
hand a  
all ex



agents which are most-instrumental  
in the production of the disease.  
I need hardly mention that wearing  
flannel next the skin especially  
during the winter season is of  
great importance.

The diet should be of the most  
simple, and abstemious kind, such  
as barley water, rice water, currant-  
jelly, dissolved in water, and other  
bland articles. Animal food in  
all cases should be strictly forbidden.



Ch

115

acute Rheumatism

No 27 Samson St.

Willie Jones

An inaugural oration

Read March 22<sup>d</sup> 1825

Acute Rheumatism

By Willie Jones of North Carolina